

NOTE

Photoinhibition in benthic diatom assemblages under light stress

P. Cartaxana^{1,*}, N. Domingues^{1,2}, S. Cruz³, B. Jesus^{2,4}, M. Laviale³, J. Serôdio³, J. Marques da Silva²

¹Centro de Oceanografia, Faculdade de Ciências da Universidade de Lisboa, 1749-016 Lisboa, Portugal

²Departamento de Biologia Vegetal and Centro de Biodiversidade, Genómica Integrativa e Funcional (BioFIG), Faculdade de Ciências da Universidade de Lisboa, 1749-016 Lisboa, Portugal

³Departamento de Biologia and CESAM – Centro de Estudos do Ambiente e do Mar, Universidade de Aveiro, Campus de Santiago, 3810-193 Aveiro, Portugal

⁴LUNAM université, Université de Nantes, Mer Molécules Santé EA 2160, Faculté des Sciences et des Techniques, BP 92208, 44322 Nantes cedex 3, France

ABSTRACT: Microphytobenthos are frequently subjected to light intensities higher than those required to saturate photosynthesis, which consequently can cause photoinhibition. Photosystem II (PSII) protein D1 (the main target of photoinhibition) and xanthophyll cycle pigments were quantified in epipelagic benthic diatom assemblages under high irradiance, in the presence of inhibitors and promoters of photoprotection mechanisms. Levels of D1 protein were significantly lower under high irradiance (1 h, 1500 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$) than under low light (80 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$), corresponding to a photoinhibition of 22 to 29%. Photoinhibition increased to 44 and 80% in the presence of lincomycin (inhibitor of chloroplast-protein synthesis) and dithiothreitol (inhibitor of the xanthophyll cycle), respectively. High light treatment had no significant effect on D1 protein concentrations in the presence of added glutathione and ascorbate, scavengers of reactive oxygen species (ROS). In contrast, the ROS promoter methylviologen increased photoinhibition to 63%. Under light stress, the functional stability of PSII reaction centres of the studied epipelagic benthic diatoms was more dependent on xanthophyll cycle activation than on D1 protein recycling mechanisms, and our results substantiate the role of antioxidants in photoprotection via ROS scavenging.

KEY WORDS: Microphytobenthos · Xanthophyll cycle · D1 protein · Reactive oxygen species · Photoprotection · Antioxidants

— Resale or republication not permitted without written consent of the publisher —

INTRODUCTION

Diatoms have developed several photoprotective mechanisms to cope with high light and minimize photoinhibition. The main physiological process involved in photoprotection in diatoms is the thermal dissipation of harmful excess energy through the diadinoxanthin (Ddx) cycle (Olaizola et al.

1994, Lavaud et al. 2004). Pigment conversion of Ddx to diatoxanthin (Dtx) induces non-photochemical quenching (NPQ) in the antenna pigment-protein complexes that decrease the excitation rate of Photosystem (PS) II reaction centres (Müller et al. 2001). Diatoms have been shown to form NPQ 3 to 5 times larger than in higher plants (Ruban et al. 2004).

*Email: pcartaxana@fc.ul.pt

Additional photoprotection mechanisms include photorespiration, cyclic electron transport and photo-reduction of molecular oxygen by PSI (see review by Niyogi 1999). The latter mechanism, associated with the scavenging of damaging reactive oxygen species (ROS) via the water–water cycle (see Asada 2006), has been shown to play an important role in the photoprotection of a planktonic diatom (Waring et al. 2010). Effective antioxidant systems employing several molecules (e.g. tocopherols, ascorbate and glutathione) and scavenging enzymes limit ROS lifetime and accumulation in the chloroplasts (Niyogi 1999, Foyer et al. 2006).

If the above-mentioned photoprotective mechanisms are insufficient to counteract overexcitation of PSII and accumulation of ROS, damage to the photosynthetic apparatus occurs. This damage is particularly relevant at the level of the PSII reaction centre protein D1 (Ohad et al. 1990, Aro et al. 1993). Fast turnover of multiple PSII subunits, including D1 protein, and the induction of transcriptional processes involved in the protection of cellular structures have been shown in diatoms at an early phase of high light exposure (Nymark et al. 2009, Wu et al. 2011). If photoinactivation exceeds the rate of repair, then photoinhibition of photosynthesis occurs because the pool of D1 protein and active PSII reaction centres decline.

Benthic microalgae (microphytobenthos) have been shown to play key roles in the productivity, trophic dynamics and sediment stability of shallow coastal ecosystems (MacIntyre et al. 1996). In intertidal sand and mudflats, diatom-dominated communities form dense surface biofilms despite the extreme variability in various key environmental parameters, including fluctuating light regimes and punctuated exposure to high light levels. Photoinhibition of PSII in microphytobenthos has been typically assessed indirectly by chlorophyll fluorescence techniques (e.g. Kromkamp et al. 1998, Perkins et al. 2010, Serôdio et al. 2012). After a light stress, the slow recovery of the quantum yield of PSII is ascribed to the photoinhibitory quenching component of NPQ, qI. However, this parameter is poorly characterized and might be due to a combination of photoprotection and photodamage (Niyogi 1999 and references therein, Müller et al. 2001).

In this study, we directly assessed photoinhibition by quantifying D1 protein immunochemically in suspensions of epipellic benthic diatoms subjected to light stress. Specific inhibitors were used to determine the relevance of the xanthophyll cycle, D1 protein recycling and ROS scavenging as photoprotection mechanisms.

MATERIALS AND METHODS

Sampling and diatom suspensions

Intact sediment cores (8 cm diameter) were collected in July (Expt 1) and September (Expt 2) 2012 during low tide from Trancão intertidal mud flats (38° 47' 46" N, 09° 05' 33" W), located in the Tagus Estuary, Portugal. Samples were brought to the laboratory and exposed to low light (80 $\mu\text{mol photons m}^{-2} \text{ s}^{-1}$) provided by a halogen lamp through fiberoptics 460-F (Walz). The epipellic fraction of the microphytobenthos community was collected using the lens tissue method (Eaton & Moss 1966). Two pieces of lens tissue (Whatman International) were placed on the surface of the sediment, and after ca. 1 h, the upper piece of lens tissue was removed and cells resuspended in filtered site water. Microscopic examination revealed that on both occasions, the epipellic community was composed exclusively of diatoms, dominated by specimens of the genera *Navicula*. Other genera represented in the microphytobenthos community included *Staurophora*, *Cylindrotheca*, *Gyrosigma*, *Nitzschia*, *Plagiotropis* and *Entomoneis*. Chlorophyll *a* concentrations in the diatom suspensions obtained ranged between 0.8 and 1.5 $\mu\text{g ml}^{-1}$.

Light and chemical treatments

Expt 1

Two different light treatments of 1 h at 20°C were applied to the diatom suspensions: low light (LL, 80 $\mu\text{mol photons m}^{-2} \text{ s}^{-1}$) or high light (HL, 1500 $\mu\text{mol photons m}^{-2} \text{ s}^{-1}$) using a temperature-controlled chamber (Fytoscope FS130, Photon Systems Instruments). The Fytoscope light source is composed of LED panels emitting in the visible and near infrared (400–775 nm). Diatom suspensions subjected to HL stress were previously subjected to 15 min of the following chemical treatments: Control (no addition); lincomycin (LINC, 0.9 mM); dithiothreitol (DTT, 2 mM); and DTT+LINC. LINC is an inhibitor of chloroplast protein synthesis, thereby preventing D1 protein recycling, and does not have short-term side effects in the concentrations used (Campbell & Tyystjärvi 2012). DTT is an inhibitor of the functioning of the xanthophyll cycle (Demmig-Adams & Adams 1993, Grouneva et al. 2008). After chemical and light treatments, diatom suspensions were rapidly filtered using syringes, filter holders and 25 mm GF/F What-

man filters. For each diatom suspension, 20 ml were filtered for protein D1 quantification and 5 ml for pigment analysis. Filtration was done under the light source of the Fytoscope, and filters were immediately frozen in liquid nitrogen.

Expt 2

LL and HL treatments were applied as described above. Diatom suspensions exposed to HL stress were previously subjected to 15 min of the following chemical treatments: Control (no addition); glutathione (GSH, 5 mM) + ascorbate (AsA, 2.5 mM); and methylviologen (MV, 1 μ M). GSH and AsA are scavengers of ROS (e.g. Noctor & Foyer 1998), while MV promotes the formation of ROS species (e.g. Mano et al. 2001 and references therein). Diatom suspensions were filtered and frozen as described above.

Protein extraction and immunodetection of D1

Diatoms were scratched off the filters into Eppendorf tubes containing 1 ml of extraction buffer (sodium phosphate 10 mM, pH 7.4; EDTA 1 mM; 0.2% Tween 20, v/v) supplemented with freshly prepared phenylmethylsulphonyl fluoride (1 mM) and DTT (2 mM). Extracts were homogenized with a vortex and frozen in liquid nitrogen. Samples were then incubated at 80°C for 5 min, sonicated (Branson 220, Branson) for 1 min and vortexed. The freeze-thaw cycle was repeated 4 times. To eliminate cell debris, the samples were centrifuged at $10\,000 \times g$ (20 min at 4°C). Protein concentrations were determined with Bradford microassay (Bio-Rad) using bovine serum albumin (Sigma-Aldrich) as a standard.

For each replicate, 2 μ g of total proteins were separated by SDS-PAGE in a 12% polyacrylamide gel using the mini-protean 3 system from Bio-Rad. Different amounts of purified PsbA/D1 protein (Agrisera) were also loaded in the gel in order to build a calibration curve to determine D1 concentrations. A protein standard (Novex, Life Technologies) was used to calculate protein size and control migration. Protein transfer to nitrocellulose membrane was performed in a Trans-Blot Semi-Dry Electrophoretic Transfer Cell (Bio-Rad) for 1 h at 140 mA in buffer (25 mM Tris, 192 mM glycine, 20% v/v methanol, pH 8.3). Protein loading was checked by Ponceau S staining. Membranes were blocked in PBS-T buffer supplemented with defatted powdered milk (5%

w/v) for 1 h at room temperature. A rabbit anti-PsbA antibody (Agrisera) was used for detection of D1, at 1:20 000 dilution in blocking buffer for 1 h. Horseradish peroxidase (GE Healthcare) coupled secondary anti-rabbit IgG antibody was incubated for 1 h at a dilution of 1:40 000. Four 15 min washings with PBS-T were performed after antibody incubation. Chemoluminescence detection was done using an ECL Advance Western Blotting Detection Kit (GE Healthcare). Developed films (Hyperfilm ECL, GE Healthcare) were imaged with a Gel Doc XR imaging system to quantify band intensities by densitometry, using Quantity-One software (all Bio-Rad). Photoinhibition was calculated as $(1 - D1_T / D1_{LL}) \times 100$, where $D1_T$ is the concentration of D1 at the different light/chemical treatments and $D1_{LL}$ is the concentration of D1 at low light.

Pigment analysis

Filters for pigment analysis were homogenized in 95% cold buffered methanol (2% ammonium acetate) using a glass rod. Samples were then sonicated for 30 s, briefly vortexed and transferred to -20°C for 30 min. Supernatants were collected after centrifugation at $1100 \times g$ (5 min at 4°C), and filtered through 0.2 μ m Fluoropore membrane filters (Millipore). Extracts were then immediately injected into an HPLC system (Shimadzu) with a photodiode array detector (SPD-M10AVP). Chromatographic separation was carried out using a Supelcosil C18 column (25 cm length; 4.6 mm diameter; 5 μ m particles; Sigma-Aldrich) for reverse-phase chromatography and a 35 min elution programme. The solvent gradient followed Kraay et al. (1992), with an injection volume of 100 μ l and a flow rate of 0.6 ml min⁻¹. Pigments were identified from absorbance spectra and retention times, and their concentrations were obtained from the signals in the photodiode array detector. Calibration curves were performed using pure crystalline standards from DHI. The de-epoxidation state (DES) was calculated as $DES = Dtx / (Ddx + Dtx)$, where Dtx and Ddx are the concentrations of diatoxanthin and diadinoxanthin, respectively.

Statistical analysis

The existence of significant differences was tested using one-way analysis of variance (ANOVA) for effects of light/chemical treatments, using in all cases (DES and D1 protein levels) 4 replicate suspensions

for each treatment. Data were transformed whenever necessary to comply with ANOVA assumptions. Post hoc comparisons were made with Tukey HSD tests. Statistical analyses were carried out using Statistica 10.0 (StatSoft).

RESULTS AND DISCUSSION

Exposure of diatom suspensions to an irradiance of $1500 \mu\text{mol photons m}^{-2} \text{s}^{-1}$ (HL) caused significantly (Tukey HSD, $p < 0.01$) lower D1 protein levels (74.1 ± 5.7 [SD] and $59.6 \pm 5.6 \text{ fmol } \mu\text{g}^{-1}$ protein) relative to LL (94.6 ± 3.6 and $84.4 \pm 5.5 \text{ fmol } \mu\text{g}^{-1}$ protein), corresponding to 22 % (Expt 1) and 29 % (Expt 2) of photoinhibition (see Figs. 1 & 2, respectively). Addition of LINC inhibited D1 protein recycling, causing a significant (Tukey HSD, $p < 0.001$) decrease in the concentrations of D1 ($52.7 \pm 4.0 \text{ fmol } \mu\text{g}^{-1}$ protein; Fig. 1). Photoinhibition was increased from 22 to 44 % in the presence of LINC.

Most phototrophs have developed a rapid D1 turnover repair cycle, involving proteolytic release of the damaged D1, de novo synthesis and incorporation of the protein into reassembled PSII complexes (Mattoo et al. 1984, Aro et al. 1993). However, significant photoinactivation by net loss of D1 protein can be attained by blocking chloroplast protein synthesis with LINC (Campbell & Tyystjärvi 2012). Recently, Domingues et al. (2012) reported significant degradation and re-synthesis of D1 protein under low irra-

diances in the diatom model species *Phaeodactylum tricornutum*, but increased turnover rates under high light. Fast turnover of multiple PSII subunits, including D1 protein, has been shown to play an important role in the resistance of small planktonic diatoms to high irradiances (Wu et al. 2011, 2012).

When exposed to DTT, an inhibitor of the xanthophyll cycle, D1 protein levels were significantly (Tukey HSD, $p < 0.001$) lower ($18.6 \pm 5.8 \text{ fmol } \mu\text{g}^{-1}$ protein) than in the presence of LINC or under LL, corresponding to a photoinhibition of 80 % (Fig. 1). No significant differences were found for DES between LL and HL treatments in the presence of DTT, showing full inhibition by DTT of Ddx conversion to Dtx. DES was significantly (Tukey HSD, $p < 0.001$) lower in LL than in both HL and HL+LINC treatments, showing activation of the xanthophyll cycle caused by HL exposure (Fig. 1). The lowest D1 protein levels ($8.9 \pm 1.7 \text{ fmol } \mu\text{g}^{-1}$ protein) were observed when both inhibitors (LINC and DTT) were used simultaneously (Fig. 1), corresponding to a photoinhibition of 91 %.

These results indicate that the functional stability of PS II in the studied epipelagic benthic diatoms under high irradiances was more dependent on xanthophyll cycle activation than on D1 protein recycling. Previous works have shown the key role of the Ddx cycle in the photoprotection of planktonic diatoms through thermal dissipation of harmful excess energy (e.g. Olaizola et al. 1994, Lavaud et al. 2004, Ruban et al. 2004). While some studies have shown that intertidal microphytobenthos depend greatly on the xanthophyll cycle during daylight emersion periods (Chevalier et al. 2010, Cartaxana et al. 2011), others estimated its contribution for photoprotection as minimal (Perkins et al. 2010, Serôdio et al. 2012).

D1 protein concentrations were not significantly different between LL and HL treatment with the addition of GSH and AsA ($82.2 \pm 10.1 \text{ fmol } \mu\text{g}^{-1}$ protein; Fig. 2). DES was significantly (Tukey HSD, $p < 0.001$) lower in LL than in all HL treatments, showing activation of the xanthophyll cycle upon exposure to high irradiances (Fig. 2). These results show full protection of PSII protein D1 under high irradiances through a combination of energy dissipation by the Ddx cycle and ROS scavenging by the added antioxidants GSH and AsA. In the chloroplasts, GSH and AsA are capable of directly quenching several ROS species (singlet oxygen, $^1\text{O}_2$; superoxide anion radical, $\text{O}_2^{\cdot-}$; and hydroxyl radical, $\text{OH}^\bullet$) and are involved in the regeneration of other antioxidant molecules (e.g. α -tocopherol; Noctor & Foyer 1998, Niyogi 1999, Foyer et al. 2006).




Fig. 1. Photosystem II protein D1 content ($\text{fmol } \mu\text{g}^{-1}$ protein) and de-epoxidation state (DES) in suspensions of benthic diatoms (mean $\pm$ SD, $n = 4$) in Expt 1. LL: low light, $80 \mu\text{mol photons m}^{-2} \text{s}^{-1}$; HL: high light, $1500 \mu\text{mol photons m}^{-2} \text{s}^{-1}$; LINC: lincomycin; DTT: dithiothreitol; Dtx: diatoxanthin; Ddx: diadinoxanthin. Different letters indicate significant differences between treatments ($p < 0.01$)




Fig. 2. Photosystem II protein D1 content (fmol μg^{-1} protein) and de-epoxidation state (DES) in suspensions of benthic diatoms (mean $\pm$ SD, $n = 4$) in Expt 2. LL: low light, 80 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$; HL: high light, 1500 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$; GSH+AsA: glutathione + ascorbate; MV: methylviologen; Dtx: diatoxanthin; Ddx: diadinoxanthin. Different letters indicate significant differences between treatments ($p < 0.01$)

Diatoms can photoreduce molecular oxygen using the Mehler reaction driven by PSI, thereby protecting PSII (Waring et al. 2010). This is an effective photoprotection mechanism provided the cells can deal with the ROS produced by a set of reactions that comprise the water–water cycle (Asada 2006). In this cycle, O_2^- is rapidly converted to hydrogen peroxide (H_2O_2) by superoxide dismutase (SOD) and reduced to H_2O by AsA, in a reaction catalysed by ascorbate peroxidase (APX). AsA is regenerated with the involvement of GSH and a set of enzymes comprising the GSH-AsA cycle. Waring et al. (2010) showed higher rates of oxygen photoreduction and increased SOD and APX activities at saturating light levels in the planktonic *Thalassiosira pseudonana* when compared to the benthic *Nitzschia epithemioides*.

Addition of MV functioned in the opposite direction, significantly (Tukey HSD, $p < 0.001$) promoting D1 protein degradation (31.5 ± 5.4 fmol μg^{-1} protein) and increasing photoinhibition under HL from 29 to 63% (Fig. 2). MV accelerates the photoproduction of O_2^- in PSI and simultaneously inhibits the photoreduction of monodehydroascorbate to AsA, inactivating APX (Mano et al. 2001). This leads to the accumulation of H_2O_2 and amplified oxidative damage.

Most available data on D1 recycling in diatoms is for planktonic, mostly centric species (e.g. Key et al. 2010, Wu et al. 2011, 2012). Studying a panel of 7 centric marine diatoms with a cell volume span of 10^1 to $10^7 \mu\text{m}^3$, Key et al. (2010) found that D1 per total protein increased with cell size and that larger spe-

cies showed slower repair rates for PSII. Further work is needed on protein D1 quantification in benthic epipellic diatoms to better characterize their response to light stress. However, a comparison with the results presented by Key et al. (2010) suggests that the species that compose the studied microphytobenthos community may allocate a high proportion of their total protein to D1 and have slow repair rates, similarly to larger planktonic diatoms.

The response of natural microphytobenthos communities to light stress is probably more complex than that of planktonic communities, due to the capacity of benthic epipellic species to migrate vertically within the sediment profile. When exposed to high irradiances, intertidal microphytobenthos have been found to migrate downwards, avoiding photoinhibitory light levels (e.g. Kromkamp et al. 1998). However, the photoprotective nature frequently attributed to this photophobic behaviour has only recently been addressed experimentally using a diatom motility inhibitor to obtain intact non-migratory biofilms (Cartaxana & Serôdio 2008, Perkins et al. 2010, Cartaxana et al. 2011). We are currently extending D1 protein quantification to sediment samples to directly assess photo-oxidative damage in intact microphytobenthos biofilms and further characterize the relevance of migration as a 'behavioural' photoprotection mechanism.

Acknowledgements. This work was supported by project MigROS (PTDC/MAR/112473/2009) funded by the Portuguese Foundation for Science and Technology.

LITERATURE CITED

- Aro EM, Virgin I, Andersson B (1993) Photoinhibition of photosystem II: inactivation, protein damage and turnover. *Biochim Biophys Acta* 1143:113–134
- Asada K (2006) Production and scavenging of reactive oxygen species in chloroplasts and their function. *Plant Physiol* 141:391–396
- Campbell DA, Tyystjärvi E (2012) Parameterization of photosystem II photoinactivation and repair. *Biochim Biophys Acta* 1817:258–265
- Cartaxana P, Serôdio J (2008) Inhibiting diatom motility: a new tool for the study of the photophysiology of intertidal microphytobenthic biofilms. *Limnol Oceanogr Methods* 6:466–476
- Cartaxana P, Ruivo M, Hubas C, Davidson I, Serôdio J, Jesus B (2011) Physiological versus behavioral photoprotection in intertidal epipellic and epipsammic benthic diatom communities. *J Exp Mar Biol Ecol* 405:120–127
- Chevalier EM, Gévaert F, Créach A (2010) In situ photosynthetic activity and xanthophyll cycle development of undisturbed microphytobenthos in an intertidal mudflat. *J Exp Mar Biol Ecol* 385:44–49

- Demming-Adams B, Adams WW III (1993) The xanthophyll cycle, protein turnover, and the high light tolerance of sun-acclimated leaves. *Plant Physiol* 103:1413–1420
- Domingues N, Matos AR, Marques da Silva J, Cartaxana P (2012) Response of the diatom *Phaeodactylum tricornutum* to photooxidative stress resulting from high light exposure. *PLoS ONE* 7:e38162
- Eaton J, Moss B (1966) The estimation of numbers and pigment content of epipelagic algal populations. *Limnol Oceanogr* 11:584–595
- Foyer CH, Trebst A, Noctor G (2006) Signaling and integration of defense functions of tocopherol, ascorbate and glutathione. In: Demmig-Adams B, Adams WW III, Mattoo AK (eds) *Photoprotection, photoinhibition, gene regulation, and environment*. Springer, Dordrecht, p 241–268
- Grouneva I, Jakob T, Wilhelm C, Goss R (2008) A new multi-component NPQ mechanism in the diatom *Cyclotella meneghiniana*. *Plant Cell Physiol* 49:1217–1225
- Key T, McCarthy A, Campbell DA, Six C, Roy S, Finkel ZV (2010) Cell size trade-offs govern light exploitation strategies in marine phytoplankton. *Environ Microbiol* 12:95–104
- Kraay GW, Zapata M, Veldhuis M (1992) Separation of chlorophylls c_1 , c_2 , and c_3 of marine phytoplankton by reversed-phase C18 high-performance liquid chromatography. *J Phycol* 28:708–712
- Kromkamp J, Barranguet C, Peene J (1998) Determination of microphytobenthos PSII quantum efficiency and photosynthetic activity by means of variable chlorophyll fluorescence. *Mar Ecol Prog Ser* 162:45–55
- Lavaud J, Rousseau B, Etienne AL (2004) General features of photoprotection by energy dissipation in planktonic diatoms (Bacillariophyceae). *J Phycol* 40:130–137
- MacIntyre HL, Geider RJ, Miller DC (1996) Microphytobenthos: the ecological role of the 'secret garden' of unvegetated, shallow-water marine habitats. I. Distribution, abundance and primary production. *Estuaries* 19: 186–201
- Mano J, Ohno C, Domae Y, Asada K (2001) Chloroplastic ascorbate peroxidase is the primary target of methylviologen-induced photooxidative stress in spinach leaves: its relevance to monodehydroascorbate radical detected with in vivo ESR. *Biochim Biophys Acta* 1504:275–287
- Mattoo AK, Hoffman-Falk H, Marder JB, Edelman M (1984) Regulation of protein metabolism: coupling of photosynthetic electron transport to in vivo degradation of the rapidly metabolized 32-kilodalton protein of chloroplast membranes. *Proc Natl Acad Sci USA* 81:1380–1384
- Müller P, Li XP, Niyogi K (2001) Non-photochemical quenching. A response to excess light energy. *Plant Physiol* 125:1558–1566
- Niyogi KK (1999) Photoprotection revisited: genetic and molecular approaches. *Annu Rev Plant Physiol Plant Mol Biol* 50:333–359
- Noctor G, Foyer CH (1998) Ascorbate and glutathione: keeping active oxygen under control. *Annu Rev Plant Physiol Plant Mol Biol* 49:249–279
- Nymark M, Valle KC, Brembu T, Hancke K and others (2009) An integrated analysis of molecular acclimation to high light in the marine diatom *Phaeodactylum tricornutum*. *PLoS ONE* 4:e7743
- Ohad I, Adir N, Koike H, Kyle DJ, Inoue Y (1990) Mechanism of photoinhibition in vivo. A reversible light induced conformational change of reaction center II is related to an irreversible modification of the D1 protein. *J Biol Chem* 265:1972–1979
- Olaizola M, Roche J, Kolber Z, Falkowski PG (1994) Non-photochemical fluorescence quenching and the diadinoxanthin cycle in a marine diatom. *Photosynth Res* 41: 357–370
- Perkins RG, Lavaud J, Seródio J, Mouget JL and others (2010) Vertical cell movement is a primary response of intertidal benthic biofilms to increasing light dose. *Mar Ecol Prog Ser* 416:93–103
- Ruban A, Lavaud J, Rousseau B, Guglielmi G, Horton P, Etienne AL (2004) The super-excess energy dissipation in diatom algae: comparative analysis with higher plants. *Photosynth Res* 82:165–175
- Seródio J, Ezequiel J, Barnett A, Mouget JL, Méléder V, Laviale M, Lavaud J (2012) Efficiency of photoprotection in microphytobenthos: role of vertical migration and the xanthophyll cycle against photoinhibition. *Aquat Microb Ecol* 67:161–175
- Waring J, Klenell M, Bechtold U, Underwood GJC, Baker NR (2010) Light-induced responses of oxygen photo-reduction, reactive oxygen species production and scavenging in two diatom species. *J Phycol* 46: 1206–1217
- Wu H, Cockshutt AM, McCarthy A, Campbell DA (2011) Distinctive photosystem II photoinactivation and protein dynamics in marine diatoms. *Plant Physiol* 156: 2184–2195
- Wu H, Roy S, Alami M, Green BR, Campbell DA (2012) Photosystem II photoinactivation, repair, and protection in marine centric diatoms. *Plant Physiol* 160:464–476